



# **Chimera States in Ensembles of Nonlocally Coupled Chaotic Systems. Control and Synchronization**

**Vadim Anishchenko**  
Saratov State University,  
Saratov, Russia


**VIII Международная научная конференция  
"КОМПЬЮТЕРНЫЕ НАУКИ И ИНФОРМАЦИОННЫЕ ТЕХНОЛОГИИ"  
памяти А.М.Богомолова**

**Саратов, 3 июля 2018 г.**

# CONTENTS

- 1. Introduction**
- 2. Basic models for analyzing chimera states in ensembles of coupled chaotic discrete-time systems**
- 3. Dynamics of an ensemble of nonlocally coupled Henon maps. Amplitude and phase chimeras**
- 4. Dynamics of a ring of nonlocally coupled Lozi maps. Solitary states**
- 5. Intermittent chimera in the Henon maps ensemble. Lifetime of amplitude chimeras**
- 6. Two coupled rings of Henon and Lozi maps. New type of chimera states. Controlling the intermittent process duration.**
- 7. Synchronization of spatio-temporal structures in coupled ensembles.**
- 8. Conclusion and publications**

# 1. Introduction. Technological Complex Networks




Traditionally the analysis of **the Internet structure** is made by means of ***traceroutes***. That is to say, by exploring all the paths from a given point to all the possible destinations.

G. Caldarelli  
CNR-INFM Centre SMC Dep. Physics University  
"Sapienza" Rome, Italy




**Power grid network** in the North America

# Complex Social Networks




**Friendship links in a school in the United States** (from G. Caldarelli)




**Knowledge Networks:** networks to capture the collective knowledge of the communities of users of online resources, such as the scientific literature and digital libraries, Wikipedia, as well as social media such as Twitter and Instagram.



# Complex Networks in Nature



**Complex structure of neuron links in the human brain**



**Figure 1.** Agent based brain model. Each of 90 gray matter brain regions is represented by a node. Lines indicate functional connections, defined by correlated functional activity measured using fMRI. Nodes may either be on (green) or off (yellow) and connections indicate positive (red) or negative (black) correlations.

# Numerical observation of chimera states: Network of nonlocally coupled identical systems

$$\dot{x}_i(t) = f(x_i(t)) + \frac{\sigma}{2P} \sum_{j=i-P}^{i+P} [f(x_j(t)) - f(x_i(t))]$$

$x_i$  are real dynamic variables ( $i = 1, \dots, N, N \gg 1$  and the index  $i$  is periodic *mod*  $N$ );

$t$  denotes time;



$\sigma$  is the coupling strength;

$P$  specifies the number of neighbors in each direction coupled with the  $i$ th element;


$r$  is the coupling radius (range),  $r = P/N$ ;

$f(x)$  defines dynamics of individual element:

- Stuart-Landau harmonic self-sustained oscillators;
- discrete-time systems (maps);
- continuous-time chaotic models;
- FitzHugh-Nagumo neural systems;
- Van der Pol oscillators;
- quantum oscillator systems;
- .....



**P=1**




**P=N/2**

# Chimera states in networks of nonlocally coupled identical oscillators


Kuramoto phase oscillator model:

$$\frac{\partial \psi}{\partial t} = \omega - \int_{-\pi}^{\pi} G(x-x') \sin[\psi(x,t) - \psi(x',t) + \alpha] dx' \quad \text{with} \quad \underbrace{G_{\text{exp}}(x) = \frac{\kappa}{2} e^{-\kappa|x|}}_{\text{Exponential coupling function}}$$

Snapshot of chimera state



average frequency



Coherent domains of periodic in-phase oscillations coexist with incoherent domains, characterized by a chaotic behavior in time and in space.

## Chimera States for Coupled Oscillators

Daniel M. Abrams\* and Steven H. Strogatz†

*Department of Theoretical and Applied Mechanics, Cornell University, 212 Kimball Hall, Ithaca, New York 14853-1503, USA*

**A chimera state** was defined as a **spatio-temporal pattern** in which an array of identical oscillators is split into **coexisting regions** of **coherent** (phase and frequency locked) and **incoherent** (drifting) oscillations.

“In Greek mythology, **the chimera** was a fire-breathing monster having a lion’s head, a goat’s body, and a serpent’s tail.

Today the word refers to anything composed of incongruent parts, or anything that seems fantastic.”



**Chimera of Arezzo**



## 2. Basic models of chaotic systems with different types of chaotic attractors

It has been shown that chimera states can be obtained only in networks of chaotic systems with **non-hyperbolic attractors** and cannot be found in networks of chaotic systems with **hyperbolic (singular-hyperbolic)** attractors.

**N. Semenova, A. Zakharova, E. Schöll, V. S. Anishchenko, Europhys. Lett. 112 (2015) 40002.**

Discrete-time system

with a non-hyperbolic attractor:

### Henon map

$$\begin{aligned}x_{n+1} &= 1 - \alpha x_n^2 + y_n \\ y_{n+1} &= \beta x_n\end{aligned}$$

It describes the properties of chaotic attractors in the Poincare section for the spiral chaos systems: the Rössler oscillator, the Anishchenko-Astakhov oscillator etc.

Discrete-time system

with a singular-hyperbolic attractor:


### Lozi map

$$\begin{aligned}x_{n+1} &= 1 - \alpha|x_n| + y_n \\ y_{n+1} &= \beta x_n\end{aligned}$$




It represents the properties of the Lorenz-type attractors in the Poincare section.

# Peculiarities of chaotic attractors

## Henon map



## Lozi map




# Ensembles of nonlocally coupled

## Henon and Lozi maps




$$x_i^{t+1} = f(x_i^t, y_i^t) + \frac{\sigma}{2P} \sum_{j=i-P}^{i+P} [f(x_j^t, y_j^t) - f(x_i^t, y_i^t)],$$

$$y_i^{t+1} = \beta x_i^t.$$





- $x_i, y_i$  are real dynamical variables ( $i = 1, 2, \dots, N$ ,  $N = 1000$  is the number of the network elements, the index  $i$  is periodic *mod*  $N$ ),
- $t$  denotes the discrete time,
- $f(x_i, y_i) = 1 - \alpha(x_i)^2 + y_i$  (Henon map) and  $f(x_i, y_i) = 1 - \alpha|x_i| + y_i$  (Lozi map),
- $\sigma$  is the coupling strength,
- $P$  specifies the number of neighbors on the left and right of the  $i$ th element,
- $\alpha = 1.4, \beta = 0.3$  are the parameters of the ensemble elements, which correspond to the chaotic dynamics in an uncoupled map,
- $r$  is the coupling radius (range),  $r = P/N$ ,
- Initial conditions are randomly distributed in the interval  $[-0.5; 0.5]$ .

# Phase and amplitude chimera states in the ensemble of nonlocally coupled Henon maps



Phase chimera for  $\sigma = 0.296$





Phase and amplitude chimeras for  $\sigma = 0.26$

I. Omelchenko, Y. Maistrenko, P. Hövel, and E. Schöll, Phys. Rev. Lett. **106**, 234102 (2011).

N.I. Semenova, E.V. Rybalova, G.I. Strelkova, V.S. Anishchenko, Regular and Chaotic Dynamics **22**(2), 148-162 (2017).



# Phase chimera



Elements in regions 1 and 2 demonstrate **weakly chaotic oscillations which are close to the 2-cycle and shifted in phase by one iteration.**

**Phase chimera** : random switchings between the in-phase (region 1) and anti-phase (region 2) oscillations.


# Cross-correlation coefficient in the phase chimera state

$$\Psi_{1,i} = \frac{\langle \tilde{x}_1(t) \tilde{x}_i(t) \rangle}{\sqrt{\langle \tilde{x}_1^2(t) \rangle \langle \tilde{x}_i^2(t) \rangle}}, \quad i = 2, 3, \dots, N$$


$$\tilde{x}(t) = x(t) - \langle x(t) \rangle$$

is a fluctuation around the average value.  
The brackets mean time averaging.


The CCC in the regime of phase chimera changes between either +1 or -1 and characterizes random switchings between the in-phase (+1) and anti-phase (-1) oscillations.




# Amplitude chimera



Snapshot



Space-time profile




Cross-correlation


**Amplitude chimera state appears** when the cluster of elements ( $120 < i < 290$ ) demonstrates developed chaotic behavior.

S.A. Bogomolov, A.V. Slepnev, G.I. Strelkova, E. Schöll, V.S. Anishchenko, Commun. Nonlinear Sci. Numer. Simul. **43**, 25 (2017).


# Temporal switchings of oscillations of the network elements in an amplitude chimera. Finite lifetime of the amplitude chimera



$\sigma = 0.304,$   
 $r = 0.32$



Amplitude chimera




Phase chimera

N.I. Semenova, G.I. Strelkova, V.S. Anishchenko, and A. Zakharova, Temporal intermittency and the lifetime of chimera states in ensembles of nonlocally coupled chaotic oscillators. CHAOS **27**, 061102 (2017).







# Dynamics of the ensemble of nonlocally coupled Lozi maps



N.I. Semenova, E.V. Rybalova, G.I. Strelkova, V.S. Anishchenko, The European Phys. J. Special Topics **226**, 1857 (2017).


# Transition to incoherence through solitary states in the network of Lozi maps for $r = 0.193$



## Mechanism of solitary state emergence in the ring of Lozi maps





$$x_i^{t+1} = f(x_i^t, y_i^t) + \frac{\sigma}{2P} \sum_{j=i-P}^{i+P} [f(x_j^t, y_j^t) - f(x_i^t, y_i^t)],$$

$$y_i^{t+1} = \beta x_i^t.$$




$$x_i^{t+1} = (1-\sigma)f(x_i^t, y_i^t) + \frac{\sigma}{2P} \sum_{j=i-P}^{i+P} f(x_j^t, y_j^t),$$

$$y_i^{t+1} = \beta x_i^t.$$




## Travelling waves in the ensemble of Lozi maps





# Network of networks



In nature and technology, it is typical the connection (and thus, the creation of a network) between different networks with different inner and external couplings.


## Two coupled rings of Henon and Lozi maps: Simple model of a network of networks

$$x_i^{t+1} = f_i^t + \frac{\sigma_1}{2P} \sum_{j=i-P}^{i+P} [f_j^t - f_i^t] + \gamma F_i^t,$$

$$y_i^{t+1} = \beta x_i^t, \quad f_i^t = 1 - \alpha(x_i^t)^2 + y_i$$

$$u_i^{t+1} = g_i^t + \frac{\sigma_2}{2R} \sum_{j=i-R}^{i+R} [g_j^t - g_i^t] + \gamma G_i^t,$$

$$v_i^{t+1} = \beta u_i^t, \quad g_i^t = 1 - \alpha|x_i^t| + y_i$$



$i = 1, 2, \dots, N = 1000$  and  $\gamma$  is the coupling coefficient between the two rings.

$$F_i^t = \bar{F}_i^t(x_i^t, y_i^t, u_i^t, v_i^t) = g(u_i^t, v_i^t) - f(x_i^t, y_i^t),$$

$$G_i^t = \bar{G}_i^t(x_i^t, y_i^t, u_i^t, v_i^t) = f(x_i^t, y_i^t) - g(u_i^t, v_i^t),$$



**Dissipative coupling**



$$F_i^t = \bar{\bar{F}}_i^t(x_i^t, y_i^t, u_i^t, v_i^t) = u_i^t - x_i^t,$$

$$G_i^t = \bar{\bar{G}}_i^t(x_i^t, y_i^t, u_i^t, v_i^t) = x_i^t - u_i^t.$$





**Inertial coupling**

# Snapshots of the dynamics of the Henon and Lozi maps rings



Henon maps

Phase and amplitude chimeras are observed in both rings.



Lozi maps







$$\gamma = 0$$

$$\gamma = 0.375$$

$$\sigma_1 = 0.32, \sigma_2 = 0.15, r_1 = 0.32, r_2 = 0.19, \alpha = 1.4, \beta = 0.3$$

# Traveling waves in the Henon and Lozi maps rings

$$\sigma_1 = 0.344,$$
$$\sigma_2 = 0.22$$







Snapshots

$$\gamma = 0.0$$


Spatio-temporal profiles

$\gamma = 0.0925$   
Traveling waves in both rings

A new type of chimera - **solitary state chimera (SSC)**  
(inertial coupling  $\gamma = 0.020$ )



## Region of existence of the solitary state chimera in the ring of Henon maps





### Solitary state chimera:

- All the elements - solitary states;
- Asynchronous chaotic behavior;
- Chaotic synchronization inside the coexisting coherence domains.





# Solitary state chimera in the Henon map ensemble (dissipative coupling $\gamma = 0.105$ )





# Temporal intermittency in the network of coupled ensembles of Henon and Lozi maps



$t_1$



$t_2$




$t_3$




$\gamma = 0.019,$   
 $\sigma_1 = 0.344,$   
 $\sigma_2 = 0.23,$   
 $r_1 = 0.32,$   
 $r_2 = 0.193,$   
 $\alpha = 1.4,$   
 $\beta = 0.3$

# Time series for the 375<sup>th</sup> element of the amplitude chimera




# Lifetime of switching process and its control in two coupled ensembles




$$\gamma = 0.015$$

$$\begin{aligned} \sigma_1 &= 0.344, \\ \sigma_2 &= 0.23, \\ r_1 &= 0.32, \\ r_2 &= 0.193, \\ \alpha &= 1.4, \\ \beta &= 0.3 \end{aligned}$$




$$\gamma = 0.003$$




$$\gamma = 0.031$$

Lifetime  $> 10^8$   
and goes to  
infinity



Switchings  
between  
amplitude  
and phase  
chimeras

## Lifetime of the amplitude chimera vs. the coupling $\gamma$



The amplitude chimera lifetime (or the intermittent process duration) depends nonlinearly on the coupling coefficient  $\gamma$ .



# Mutual synchronization of spatio-temporal structures in coupled ensembles

When the parameters are varied in the system of coupled ensembles (1), identical spatial structures appear simultaneously in both ensembles.

**The question is whether these structures can be synchronized and what is the region of their synchronization?**


The identity of the observed structures is diagnosed by using **the mutual correlation coefficient**:

$$R_i = \frac{\langle \tilde{x}_i(t) \tilde{u}_i(t) \rangle}{\sqrt{\langle \tilde{x}_i^2(t) \rangle \langle \tilde{u}_i^2(t) \rangle}}, \quad \tilde{x}(t) = x(t) - \langle x(t) \rangle$$


is a fluctuation around the average value. The brackets mean time averaging.

$$i = 1, 2, \dots, N$$

$\sigma_{\text{Henon}} = 0.32$   
 $\sigma_{\text{Lozi}} = 0.15$





$\sigma_{\text{Henon}} = 0.18$   
 $\sigma_{\text{Lozi}} = 0.15$



Spatio-temporal structures in the Henon maps (left), the Lozi maps (middle) ensembles and the mutual correlation coefficient (right)


## Synchronization region



# External and mutual synchronization in a two layer network of nonlocally coupled logistic maps

$$x_i^{t+1} = f_i^t + \frac{\sigma_1}{2R} \sum_{j=i-P}^{i+P} [f_j^t - f_i^t] + \gamma_{21} F_i^t,$$

$$y_i^{t+1} = g_i^t + \frac{\sigma_2}{2P} \sum_{j=i-R}^{i+R} [g_j^t - g_i^t] + \gamma_{12} G_i^t.$$



$$f_i^t = \alpha_1 x_i^t (1 - x_i^t), \quad g_i^t = \alpha_2 y_i^t (1 - y_i^t), \quad i = 1, 2, \dots, N = 1000$$

$\alpha_1$  and  $\alpha_2$  are the control parameters which define the local dynamics;

$\sigma_1$  and  $\sigma_2$  are the nonlocal coupling strengths (intra-couplings);

$P = R = 320$  denote the number of neighbors of the  $i$ th element from each side;

$F_i^t = (g_i^t - f_i^t)$ ,  $G_i^t = (f_i^t - g_i^t)$  are the coupling functions;

$\gamma_{21}$  and  $\gamma_{12}$  are the coupling strengths between the sub-networks (inter-coupling);

Initial conditions are randomly distributed in the interval  $[0; 1]$  for both rings.

# Uncoupled case


Parameter mismatch:

$$\alpha_1 = 3.7, \alpha_2 = 3.85$$


Different intra-couplings:

$$\sigma_1 = 0.23, \sigma_2 = 0.15$$

First ring






Second ring




Phase and amplitude chimeras

Spatiotemporal chaos


# External synchronization of chimera states



$$\gamma_{12} = 0.15$$




$$\gamma_{12} = 0.3$$



$$\gamma_{12} = 0.45$$









# External synchronization of spatiotemporal patterns




$\alpha_1$  is varied in  $[3.5, 3.95]$ ,

$\alpha_2 = 3.85$ ,  $\gamma_{12} = 0.4$




## Region of external synchronization in the $(\alpha_1, \gamma)$ parameter plane




- Inside the synchronization regime, the state of the driver network **fully determines** the state in the response network.
- The structure in the response network **“is locked”** by the pattern in the driver network.

## Mutual synchronization of chimera structures



The chimera structures are synchronized at  $\gamma = 0.075$

## Region of Synchronization



# Conclusions

1. The basic models of discrete-time chaotic systems (the Henon and Lozi maps) have been selected to describe the spatiotemporal dynamics in ensembles of nonlocally coupled chaotic oscillators.
2. Possible spatiotemporal structures realized in those models have been analyzed in detail.
3. A novel type of chimera states, a solitary state chimera, has been revealed.
4. It has been shown that the amplitude chimeras are non-stationary and are characterized by a finite lifetime.
5. Effects of external and mutual synchronization of chimera states have been established and analyzed.

**This research is carried out in the framework of the SFB 910 project.**

# My co-authors

## Saratov State University



St. Sergey  
Bogomolov



Prof. Tatiana  
Vadivasova



Dr. Galina  
Strelkova



Dr. Nadezhda  
Semenova



St. Elena  
Rybalova



PhD st.  
Andrei Bukh

## Technical University of Berlin, Germany



Prof. Dr. Eckehard  
Schöll



Dr. Anna  
Zakharova



## Our papers

1. N. Semenova, A. Zakharova, E. Schöll, V. S. Anishchenko, *Europhys. Lett.* **112** (2015) 40002.
2. N. Semenova, A. Zakharova, E. Schöll, and V. Anishchenko, *AIP Conference Proceedings* 1738, 210014 (2016); doi: 10.1063/1.4951997.
3. S.A. Bogomolov, G.I. Strelkova, E. Schöll, and V.S. Anishchenko, *Amplitude and phase chimeras in an ensemble of chaotic oscillators. Tech. Phys. Lett.* **42**(7), 763-766 (2016).
4. S.A. Bogomolov, A.V. Slepnev, G.I. Strelkova, E. Schöll, and V.S. Anishchenko, *Mechanisms of Appearance of Amplitude and Phase Chimera States in Ensembles of Nonlocally Coupled Chaotic Systems. Commun. Nonlinear Sci. Numer. Simulat.* **43**, 25-36 (2017).
5. T.E. Vadivasova, G.I. Strelkova, S.A. Bogomolov, and V.S. Anishchenko, *Correlation analysis of the coherence-incoherence transition in a ring of nonlocally coupled logistic maps. CHAOS*, **26** (2016).
6. N.I. Semenova, E.V. Rybalova, G.I. Strelkova, and V.S. Anishchenko, “Coherence-incoherence” Transition in Ensembles of Nonlocally Coupled Chaotic Oscillators with Nonhyperbolic and Hyperbolic Attractors. *Regular and Chaotic Dynamics*, **22**(2), 148-162 (2017).
7. N.I. Semenova, G.I. Strelkova, V.S. Anishchenko, and A. Zakharova, *Temporal intermittency and the lifetime of chimera states in ensembles of nonlocally coupled chaotic oscillators. CHAOS*, **27**, 061102 (2017).
8. A. Bukh, E. Rybalova, N. Semenova, G. Strelkova, and V. Anishchenko, *New type of chimera and mutual synchronization of spatiotemporal structures in two coupled ensembles of nonlocally interacting chaotic maps. CHAOS*, **27**(11), 111102 (2017).

**Thank you very much  
for your attention!**